



PREVACC should be used before the Vaccinations, **PREVACC** Protects the vaccination, improving the quality of the Vaccinations.

It remove chlorine and chloramines completely. **PREVACC** has a Brilliant Blue Supra FCF colour, Which allows to see its distribution in the water system and to monitor vaccine distribution and administration in the water system for poultry, aquaculture, ect.

Unique feed supplement to achieve most effecient 'take' of live vaccine.

- Neutralize chlorine levels up to 5 PPM in water and improve vaccination effeciency.
- Contains a blue colour, vaccine administration can be monitored at Birds Break, tounge and crop.
- Substitutes the use of skimmed milk powder and therefor reduces bacterial contamination or blockages of the water system.
- Does not affect waters intake.

Store in cool dry place. Protected from direct sunlight.

Dosage:

20 ml in 100 Litres of water, wait for 10 minutes for complete neutrilization of chlorine up to 5 PPM. Stir gently to mix the solution.

How to use / Mode of use:

01. Shake well before use
02. Determine the water volume used in water vaccination
03. Add liquids as per Dosage / Requirement
04. Wait for 10 minutes to allow complete dissolution & neutralization of chlorine
05. Stir gently to mix solution. For better performance
06. Follow recommendations for application of the vaccine, medicine or probiotic

07. Add the vaccine, medicine or probiotic and mix properly

08. Even distribution of the vaccine solution in to water is indicated by the blue dye

Precautions:

Keep the product out of the reach of children. If the product gets into your eyes rinse them with abundant running water for at least 10 minutes